

PEDIATRIC	OTHER USES
PAIN MANAGEMENT	WOUND CARE
GYNEOLOGICAL / OB	WEIGHT MANAGEMENT
DERMATOLOGY	VETERINARY
DIABETES CARE	SINUS CARE
DENTAL	SCAR CARE
BIOIDENTICAL HORMONE (BHRT)	PODIATRY



CASE REPORT

— SCARS —

STRETCH MARKS PCCA PRACASIL™-PLUS



1100 Canyon View Dr, Suite C
 Santa Clara, UT 84765
 Phone: (888) 963-9923
 Fax: (855) 853-3465
 Email: rx@rx-fusion.com
 Web: www.FusionSpecialtyPharmacy.com

STRETCH MARKS

PCCA PRACASIL™-PLUS

— FUSION SPECIALTY PHARMACY —

SUMMARY: Veronica is a 34 year old female that resented with stretch marks covering her belly and thighs after pregnancy. Her use of Fusion Specialty Pharmacy's Scar Gel was for 180 days.

INTRODUCTION

Stretch marks are formed when the skin has been stretched and damaged. They are often indented streaks that are discolored with both hyperpigmented (dark colors) and hypopigmented (white) marks. Stretch marks are not painful and do not cause harm to the body, however they may be cosmetically unappealing [1].

Stretch marks usually form in areas of rapid weight gain or loss. They are common during and after pregnancy where the skin of the stomach stretches to fit around the baby or in other areas of the body experiencing rapid weight gain. Other common sites of stretch marks include the breasts, hips, buttocks, and thighs [1].

There are many home remedies thought to prevent or treat stretch marks such as cocoa butter and Vitamin E. Although these methods are not harmful, they most likely will not reduce the appearance of stretch marks. Remedies that do tend to show improvement include retinoid creams, light and laser therapy, and microdermabrasion. These methods stimulate collagen growth and new skin production which lightens the area of hyperpigmentation.

Fusion Specialty Pharmacy has developed a gel to lighten and smooth out stretch marks to make them less visible. This gel includes Collagen, Glycolic Acid, Pentoxifylline, and Tretinoin in a proprietary base.

CASE REPORT

A 34 year old female with stretch marks covering her belly and thighs after pregnancy. The clinical pharmacists from Fusion Specialty Pharmacy worked with patient's doctor to customize a prescription stretch mark gel. The patient's stretch marks began to fade in color and the indentations started to become more even and smooth. After 45 days, the stretch marks were significantly reduced. By 180 days of using the gel, her skin had significantly evened out and the stretch marks were hardly visible.

STRETCH MARKS



SCAR GEL INFORMATION: Fusion Specialty Pharmacy has developed a gel to lighten and smooth out stretch marks to make them less visible. This gel includes Collagen, Glycolic Acid, Pentoxifylline, and Tretinoin.

Collagen	0.3%
Glycolic Acid	1%
Pentoxifylline	0.5%
Tretinoin.....	0.01%
Proprietary base.....	QS

CONCLUSION

This customized prescription therapy for topical stretch mark helped promote healing and reduce the appearance of patient’s stretch marks. Topical collagen may help reduce the appearance of lines and marks on the skin; it also helps hydrate the skin, which is important for skin healing [4]. Glycolic acid stimulates removal of dead skin cells and regrowth of new skin. It has been shown to be effective in reducing hyperpigmented skin damage [5]. Pentoxifylline prevents abnormal collagen synthesis during skin healing [6]. Tretinoin is a naturally occurring vitamin A derivative that regulates cell reproduction. It is useful in treating skin blemishes and reducing hyperpigmentation [1].

As seen in this case, a customized prescription for Fusion’s stretch mark gel therapy significantly improved the appearance of this patient’s stretch marks. After continued use of this gel for 180 days, the stretch marks were barely visible. This gel allowed for the reduction of this patient’s stretch marks by simply applying a gel to the area without the use of lasers or physical intervention.



REFERENCES:

1. Korgavkar, K. and Wang, F. (2015), Stretch marks during pregnancy: a review of topical prevention. *Br J Dermatol*, 172: 606–615. doi:10.1111/bjd.13426
2. El Taieb MA, Ibrahim AK. Fractional CO2 Laser Versus Intense Pulsed Light in Treating Striae Distensae. *Indian Journal of Dermatology*. 2016;61(2):174-180. doi:10.4103/0019-5154.177774.
3. Wang, F., Calderone, K., Smith, N.R., Do, T.T., Helfrich, Y.R., Johnson, T.R.B., Kang, S., Voorhees, J.J. and Fisher, G.J. (2015), Marked disruption and aberrant regulation of elastic fibres in early striae gravidarum. *Br J Dermatol*, 173: 1420–1430. doi:10.1111/bjd.14027
4. Schwartz SR, Park J. Ingestion of BioCell Collagen®, a novel hydrolyzed chicken sternal cartilage extract; enhanced blood microcirculation and reduced facial aging signs. *Clinical Interventions in Aging*. 2012;7:267-273. doi:10.2147/CIA.S32836.
5. Erbagcı, Z. and Akçali, C. (2000), Biweekly serial glycolic acid peels vs. long-term daily use of topical low-strength glycolic acid in the treatment of atrophic acne scars. *International Journal of Dermatology*, 39: 789–794. doi:10.1046/j.1365-4362.2000.00076.x
6. Leventhal, D., Furr, M., Reiter, D. Treatment of Keloids and Hypertrophic Scars *Arch Facial Plast Surg*; 2006;

